

## How can I obtain a copy of the Iowa Satisfaction with Anesthesia Scale? For what uses is it appropriate?

To download a copy as an Acrobat (PDF) file and the required agreement for use, [click here](#). The paper describing the instrument is: [Development of a measure of patient satisfaction with monitored anesthesia care: the Iowa Satisfaction with Anesthesia Scale](#). Validation and study design for multiple centers is described in: [Multicenter assessment of the Iowa Satisfaction with Anesthesia Scale, an instrument that measures patient satisfaction with monitored anesthesia care](#). The instrument should **not** be used for general anesthesia, regional anesthesia, sedation for non-surgical procedures, or deep sedation (i.e., patient unconscious for most of the surgery, [click here](#)). For a study of the instrument for cataract surgery patients only, [click here](#). For a reliable and valid Spanish translation, [click here](#). For a French version, [click here](#). For a Turkish version, [click here](#). The scale should be administered no sooner than 1-hour after the last sedative dose, [click here](#).

For papers reviewing methods of measuring patient satisfaction with anesthesia care, [click here](#), [click here](#), and, (for the most recent) [click here](#).

For patients undergoing general anesthesia, several instruments measure patient satisfaction with anesthesia and surgery (i.e., not the anesthetic itself). For relevant papers, [click here](#), [click here](#), [click here](#), [click here](#), and [click here](#). For measuring patient satisfaction with regional anesthesia, [click here](#). For satisfaction with the perioperative period including regional anesthesia, [click here](#). For satisfaction with the ambulatory surgery experience, preanesthesia through discharge, [click here](#). For satisfaction with sedation for endoscopy, [click here](#) and [click here](#). For satisfaction with sedation and other efficacy outcome measures, [click here](#). To understand why there is not evaluation of satisfaction with individual anesthesia providers, [click here](#).

For applications on providing information to patients, [click here](#).

If your goal is to measure patients' recovery of functional ability after surgery, [click here](#).

Return to [Frequently Asked Questions](#)